



---

## Comparative Analysis of Dietary Silica in *Vigna subterranean* (Bambara Nut) and *Arachis hypogea* (Ground Nut or Peanut)

Terhemba Simeon Iningev\*, Ochanya Elizabeth Agbo, Tordue Gabriel Buluku, Otuwole Esther Abu

Department of Chemistry, Faculty of Science, Benue State University, PMB 102119 Makurdi, Nigeria

\*Corresponding author. Terhemba Simeon Iningev

\*E-mail: [iningevt@gmail.com](mailto:iningevt@gmail.com)

### Abstract

Nutritional awareness has been longtime challenge in Nigeria even among literate populace. Many Nigerians preferred food that has characteristic taste to nutritional value. A comparative study to estimate the amount of silica in bambara nut (BN) and ground nut (GN) was carried out in order to create awareness on the preferred grain to be consumed, due to nutritional health benefits of silica. The pulverized sample of each cotyledon was carbonized at 350 °C for 30 minutes using muffle furnace to obtain bambara nut ash (BNA) and ground nut ash (GNA), respectively. To produce silica, 10 g ash of each sample was digested separately with NaOH solution to form sodium trioxosilicate. The solution was treated with HCl solution to yield silica at pH adjusted to 7.0. The yield of 2.29% and 9.66%, for BN and GN, respectively, was obtained. Characterization using FT-IR technique has indicated the presence of silanol (Si-OH) and siloxane (Si-O-Si) functional groups in the spectra of the two samples. In addition, Scanning Electron Microscope (SEM) analysis respectively revealed the morphological structure of SiO<sub>2</sub> extracted from BNA and GNA to be amorphous. Thus, the form in which cells of the body can assimilate.

**Keywords:** Bambara nut, Ground nut, Silica, Characterization, FT-IR and SEM

---

### 1. Introduction

Elements like nitrogen, phosphorus and potassium are essential nutrients without which plants cannot complete their life cycle. While element like silicon (Si) is non-essential, but is of beneficial to plants and provide certain health benefits to higher animals [1-2]. Si is the second most abundant element present in the earth's crust after oxygen. Silicon dioxide or silica (SiO<sub>2</sub>); the chemically combined form of the element is the most important Si-containing inorganic substance [3]. It occurs in crystalline, poorly crystalline and amorphous phases. Amorphous silicon dioxide or silica is an important natural source that release ortho-silicic acid as a bioavailable form of silicon. Amorphous silica is acid anhydride of monomeric ortho-silicic acid (H<sub>4</sub>SiO<sub>4</sub>) that is water soluble and highly stable in diluted aqueous solution. Other hydrated forms of ortho-silicic acid include meta-silicic acid (H<sub>2</sub>SiO<sub>3</sub> or lower oligomers such as di-silicic (H<sub>2</sub>Si<sub>2</sub>O<sub>5</sub>) and tri-silicic acids (H<sub>2</sub>Si<sub>3</sub>O<sub>7</sub>), all these can exist in aqueous solutions [3-4]. Silicic acid [Si(OH)<sub>4</sub>], is an uncharged monomeric molecule at a pH of solution below 9, and it is the form in which Si is taken up by the plant roots from the soil solution. The ability of plants to absorb and accumulate Si differs considerably, ranging from 0.1 % to 10.0 % (dry weight). However, this depends much on the plant species and Si



concentration within the vicinity [5]. Researchers have reported the absorption of Si by monocotyledonous and dicotyledonous species. According to the report, monocots in their tissues, accumulate Si in the order of 5% or more on a dry weight basis and are described as Si accumulators. The dicots on the other hand contain about 0.1% Si on a dry weight basis, and are portrait as intermediate accumulators [6]. The difference in the Si uptake and accumulation of the two cotyledons is that; in the monocots, Si uptake is an active process whereas for dicotyledonous plants, Si is absorbed passively [7]. The absorption of Si by plants from soil in the form of silicic acid is accumulate as silica, where it gets deposited on epidermal cells or as phytoliths in lumen cells. Silicon in the food is ingested as silica when feed by man and other animals [2].

The ortho-silicic acid or amorphous silica is a major form of bioavailable source of silicon that plays numerous key roles in plants, animals and human's health. It has been reported that the presence of silica in plants help to alleviate many abiotic and biotic stresses, leading to protective effect such as physical barrier to penetration and reduction in enzymatic degradation by fungal and other pathogens [8]. It has also been reported that single mineral, typically calcium is not the only mineral involved in bone health formation. It worked synergistically with other minerals such as silica in bone maintenance. Silica is most important to elderly people because it help to prevent the decalcification of bones and improve bones tensile strength [9]. People having silica deficiency are prompt to be affected by diseases like Alzheimer which is caused by accumulation of aluminium in brain tissues [10].

Silica is an important optimal for collagen synthesis that activate the hydroxylation enzymes that improve skin strength and elasticity. It was suggested that hair strands with higher silica content have lower falling rate and higher brightness. Silica also improves quality of nails, and increase protection against nail infections. Deficiency of it can lead to soft and brittleness of nails because silicon is one of the predominant element in nail composition [11]. All the above mentioned problems were in connection with people that are advanced in age. Other health benefits include protection of the heart by reducing atherosclerosis risk, or lowering the cholesterol level in order to harden the arteries. The University of Memphis (2018) reported that the mineral (silica) can support the immune response and also help to control body inflammation, thereby keeping the heart in good working conditions [12]. Also another studies carried out on some animals have proved that silicon ingested in form of silica helps to prevent the occurrence of streptozotocin-induced diabetes [13].

Though despite the extra-ordinary medicinal and nutritional values one can derive from *Vigna subterranean* and *Arachis hypogea*, there are some ill-health linked with consumption of the two cotyledons. BN and GN are both contaminated by *Aspergillus* species (*A. flavus*, *A. parasiticus*, and *A. nomius*) due to poor agricultural practices (preharvest and postharvest conditions). These fungi produced aflatoxins that are known to cause cancer in humans. This significant risk is associated to eating the seeds raw. Besides, GN is also known to contain anti nutrients factors such as phytic acid, trypsin inhibitor, tannin, saponin, which inhibit absorption of some nutrients in diet [13-15]. However, we have seen the nutritional, biological and therapeutic effects of silica; it's depletion level in our body system via kidney and skin flushing needs to be replenished by eating the right meal that contain relative amount of silica. Though many literatures had reported food sources that contain silica [3,13,16-18], but published literatures on comparing the amount of silica in edible food is rare. Hence the purpose of this paper has been aimed at comparative analysis of dietary silica in *Vigna subterranean* and *Arachis hypogea*, respectively.

## 2. Materials and Methods

### Materials

Sodium hydroxide purchased from Merck, hydrochloric acid from QualiKems, and distilled water were used as received.


### Methods

BN and GN samples were washed to remove impurities. These were dried for four days and pounded separately using mortar and pestle. The pounded samples were carbonized at 350 °C for 30 minutes using a muffle furnace to obtain BNA and GNA, respectively. 10 g, each of the carbonized sample was weighed into a 250 mL separate conical flask. 40 mL 0.1 M NaOH solution was added and heated at 60 °C for 1h with constant stirring. The solution was allowed to cool to ambient temperature and filtered using Whatman filter paper No 42. The filtrate was



treated with 0.1 M HCl with stirring until the pH of the solution was adjusted to 7.0. The silica formed was aged for 24 h; filtered, washed and dried at 40 °C for 4h. It was stored in an airtight plastic bag prior for analysis.

The reaction involving conversion of silicon dioxide to sodium trioxosilicate (IV) is shown in equation (1), whereas production of silica is shown in equation (2).



Treatment of sodium trioxosilicate with HCl forms silica





Figure 1: Plates (a) BN seeds (b) GN seeds (c) Pounded BN seeds (d) Pounded GN seeds (e) Crystal formation from BNA (f) Crystal formation from GNA (g) Filtered crystal formation from BN sample (h) Filtered crystal formation from GN sample

### Determination of Physicochemical Property of Silica

#### Percentage yield

The percentage yield of the silica extracted was calculated using the formula

$$\% \text{ Yield} = \frac{\text{Actual Yield (g)}}{\text{Sample (g)}} \times 100$$

#### Characterization

##### FT-IR Measurement

IR spectrum was recorded as Nujol mulls using KBr pellets on FT-IR spectrophotometer Shimadzu (FT-IR 8400S). One milligrams (1 mg) of each sample was weighed into a small agate mortar and a drop of nujol was added and ground. The mull obtained was suspended on the cell of the spectrophotometer and scanned between 4000-400  $\text{cm}^{-1}$  at 32 runs per minute.

##### SEM Measurement

The morphology of the prepared silica powder was investigated by SEM (Quanta 400, FEI, USA). The sample was coated with a very thin film of gold (Au) using a sputter coater (SPI, 12150 AX, USA) and prior to SEM analysis.

### 3. Results and Discussions

#### Percentage Yield of Silica Extracted

The percentage yields of silica extracted from BNA and GNA digested with 0.1 M NaOH solution were found to be 2.29 % and 9.66 %, respectively (Table 1). The results indicate that silica is present in both the mono- and dicotyledon grains. From the study it was observed that GN has higher yield compare to BN. Though the literature



[6] established that monocot (Si accumulator) grains have higher silica content than dicot. However, the higher yield observed could be attributed to the oil present in GN [19]. In this study, the yield of the monocot is <5.0 %; this could be probably due to incomplete conversion of Si to SiO<sub>2</sub> during carbonization. Besides, literatures [20-21] have shown that concentration and volume of alkaline solvent as well as extraction temperature and time factors can affect the yield of silica. It means, the more is the solvent, the higher the surface area for the distribution of solvent to ash. Hence there would be increased in the formation of Na<sub>2</sub>SiO<sub>3</sub> as well as the yield of silica.

**Table 1:** Percentage yield of the extracted silica

Sample	Percentage Yield
BN	2.29
GN	9.66

### Infrared Spectral Analysis

The FT-IR spectra of silica extracted from BN and GN carbonized at 350 °C were shown in Fig. 2a and 2b, respectively. A peak at 1654.9 cm<sup>-1</sup> and a broad band that appeared at 3298.7 cm<sup>-1</sup> (Fig.2a) was ascribed to bending and stretching vibrations of the O-H bond from the silanol groups (Si-OH), respectively. This implies that water a molecule is trapped on the silica surface [22]. Similar vibration bands (1654.9 cm<sup>-1</sup> and 3208.1 cm<sup>-1</sup>) were observed on spectrum of GN (Fig.2b).




Figure 2a: FT-IR spectrum for silica (BN)





Figure 2b: FT-IR spectrum for silica (GN)



The peaks at  $1017.6\text{ cm}^{-1}$  and  $1028.7\text{ cm}^{-1}$  from Fig.2a and 2b, respectively, were ascribed to Si-O asymmetric stretching vibration [23]. Peaks at  $1148.0\text{ cm}^{-1}$  (Fig.2a) and  $1230.0\text{ cm}^{-1}$  (Fig.2b) were assigned to stretching mode of Si-O-Si (siloxane), which clearly showed silica formation. The absorption frequency bands observed at  $2922.2\text{ cm}^{-1}$  and  $2855.1\text{ cm}^{-1}$  from both figures were ascribed to bending mode of  $-\text{CH}_2$  and  $-\text{CH}_3$ , respectively. Appearance of these bands in the spectra signifies the presence of an organic compound in the two samples [24-25].

#### 4. SEM measurement

Images of silica obtained from Vigna subterranean and Arachis hypogea as well as total analysis were depicted in Fig. 3a and Fig.3b, respectively. The images have shown that the prepared silica derived from both samples carbonized at  $350\text{ }^\circ\text{C}$  were of average size particle and irregular in shape, that is, amorphous [23]. SEM analysis also showed that silicon content in BN (2.13 weight concentration) is more than in GN (1.09 weight concentration); see Fig.3a and Fig.3b. This information is in supportive of the literature which established that, monocotyledon grains are Si accumulator whereas dicotyledons are Si intermediate, as respectively shown in Fig.3a and Fig.3b.



Element Number	Element Symbol	Element Name	Atomic Conc.	Weight Conc.
6	C	Carbon	83.16	61.25
17	Cl	Chlorine	2.59	5.63
22	Ti	Titanium	1.68	4.94
13	Al	Aluminium	2.93	4.85
41	Nb	Niobium	0.77	4.38
39	Y	Yttrium	0.68	3.71
15	P	Phosphorus	1.53	2.90
16	S	Sulfur	1.43	2.81
14	Si	Silicon	1.24	2.13
19	K	Potassium	0.89	2.12
11	Na	Sodium	1.49	2.10
12	Mg	Magnesium	1.15	1.71
23	V	Vanadium	0.47	1.47
26	Fe	Iron	0.00	0.00
20	Ca	Calcium	0.00	0.00

Figure 3b: Micrograph of GN at  $750\times$  magnification

#### 5. Conclusion

The yield of silica (BN: 2.29 %, GN: 9.66 %) obtained from 10 g ash of each grain using 0.1 M NaOH analytical reagent revealed that BN and GN contain silica. The FT-IR data had shown that the prepared silica contain silanol and siloxane groups. In addition, SEM analysis depicted amorphous state of the extracted silica and has shown that, Si in BN is higher compared to GN. We therefore, recommended consumption of more BN to promote healthy bone formation, improve skin strength and elasticity, lower falling rate of hair, lowers cholesterol level thereby reducing the risk of heart disease such as atherosclerosis.

#### Acknowledgements

The authors gratefully acknowledged the National Research Institute of Chemical Technology, Zaria, Nigeria, for running the FTIR and SEM analyses. We thank technical staff headed by Dr. G.H. Atoo at the Department of Chemistry Benue State University, Makurdi-Nigeria, for their assistance during the work.



### Conflicts of Interest

Regarding the publication of this paper, the authors declare no conflicts of interest.

### References

- [1]. Uchida R. (2000). Essential Nutrients for Plant Growth: Nutrient Functions and Deficiency Symptoms. *Plant Nutrient Management in Hawaii's Soils, Approaches for Tropical and Subtropical Agriculture*. J. A. Silva and R. Uchida, eds. College of Tropical Agriculture and Human Resources, University of Hawaii at Manoa.
- [2]. Vasanthi N., Lilly M. S. and Anthoni S. R. (2012). Silicon in Day Today Life. *World Applied Sciences Journal*, 17 (11): 1425-1440.
- [3]. Jurkić L. M., Ivica C., Sandra K. P. and Krešimir P. (2013). Biological and therapeutic effects, of ortho-silicic acid and some ortho-silicic acid-releasing compounds: New perspectives for therapy. *Nutrition and Metabolism*, 10:2.
- [4]. Michael S., Danuta K., Yakov K. and Jörn B. (2006). Silicon pools and fluxes in soils and Landscapes—a review. *Journal of Plant Nutrition and Soil Sciences*, 169, 310–329.
- [5]. Jian F. Ma and Naoki Y. (2006). Silicon uptake and accumulation in higher plants. *Trends in Plant Science*, 11(8).
- [6]. Mvondo-She M. A. and Diana M. (2019). The Investigation of Silicon Localization and Accumulation in Citrus. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6681401/> Retrieved on 28/5/2021.
- [7]. Yongchao L., Jin Si and Volker R. (2005). Silicon uptake and transport is an active process in *Cucumis sativus*. *New Phytologist*, 167: 797–804.
- [8]. Heather A. C. and Carole C. P. (2007). Silica in Plants: Biological, Biochemical and Chemical Studies. *Annals of Botany*, 100(7), 1383–1389.
- [9]. Charles T. P., Kenneth J. K. and Joshua R. L. (2013). Silicon: A Review of Its Potential Role in the Prevention and Treatment of Postmenopausal Osteoporosis. *International Journal of Endocrinology*, Volume 2013, Article ID 316783, 1-6. <http://dx.doi.org/10.1155/2013/316783>
- [10]. Ambreen M., Andrew K., Claire T. and Christopher E. (2017). Aluminium in brain tissue in Familial Alzheimer's disease. *Journal of Trace Elements in Medicine and Biology*, 40, 30-36.
- [11]. Lidiane A. de A., Flavia A. and Patrícia M.B.G.M.C. (2016). Use of silicon for skin and hair care: an approach of chemical forms available and efficacy. *Anais Bras Dermatol*, 91(3): 331–335. doi: 10.1590/abd1806-4841.20163986
- [12]. Matin K.R (2018) Dietary Silicon is biofortification essential. *Journal of Nutrition and Food Science Forecast*; 1(2): 1006.
- [13]. Anna B. C. (2011). Silicon in Medicine and Therapy. *Journal of Elementology*, 16(3):489-497.
- [14]. Tan XL, Azam-Ali S., Goh E., Mustafa M., Chai HH., Ho WK., Mayes S., Mabhaudhi T. and Massawe F. (2020). Bambara Groundnut: An Underutilized Leguminous Crop for Global Food Security and Nutrition. *Frontiers in Nutrition*, 7:601496. doi:10.3389/fnut.2020.601496.
- [15]. Makinde F.M. and Dauda D.S. (2020). Nutritive Value and Inherent Anti-Nutritive Factors in Processed Peanut (*Arachis hypogaea*). *Sustainable Food Production*, 8: 17-28.
- [16]. Namiki M. and Jian F. M. (2005). Uptake system of silicon in different plant species. *Journal of Experimental Botany*, 56(414).
- [17]. Nudrat A.A., Fahad S. and Muhammad A. (2018). Peanut (*Arachis hypogaea* L.): A Prospective Legume Crop to Offer Multiple Health Benefits under Changing Climate. *Comprehensive Reviews in Food Science and Food Safety*, 17(5): 1325-1338.
- [18]. Guntamukkala B. R. and Pusarla S (2017). Silicon uptake, transportation and accumulation in Rice. *Journal of Pharmacognosy and Phytochemistry*, 6(6): 290-293.



- [19]. Yaduru S., Manish K. et al (2017). Molecular mapping of Oil Content and FattyAcids Using Dense Genetic Maps in Groundnut (*Arachishypogaea* L) front. *PlantScience* <https://doi.org/10.3389/Fp/s.2017.00794>
- [20]. Addis, L. J. (2018). Extraction of Silica Gel from Rice Husk Ash for Promising Sustainable Industrialization: Statistical Analysis. *Journal of Materials Science and Engineering with Advanced Technology*, 18(1-2): 21-42.
- [21]. Setyawan, N., Hoerudin and Wulanawati, A. (2019). Simple Extraction of Silica Nanoparticles from Rice Husk Using Technical Grade Solvent: Effect of Volume and Concentration. *IOP Conference. Series: Earth and Environmental Science*, 309: 1-8.doi:10.1088/1755-1315/309/1/012032.
- [22]. Panca S. U., Ram Y. and Chayanoot S. (2019). Production and Characterization of Precipitated Silica from Palm Oil Mill Fly Ash Using CO2 Impregnation and Mechanical IFragmentation. *Brazilian Journal of Chemical Engineering*, 36(01): 523 – 530.
- [23]. Chanadee T. and Chaiyarat S. (2016). Preparation and Characterization of Low Cost Silica Powder from Sweet Corn Cobs (*Zea mays saccharata* L.). *Journal of Material and Environmental Sciences*, 7(7): 2369-2374.
- [24]. Van H. L., Chi N. H. T. and Huy H.T. (2013). Synthesis of silica nanoparticles from Vietnamese rice husk by sol–gel method, *Nanoscale Research Letters a Springer Open Journal*, 8(58): 1-10.
- [25]. Ananthi A., Geetha D. and Ramesh P. S. (2016). Preparation and Characterization of Silica Material from Rice Husk Ash – An Economically Viable Method. *Chemistry and Materials Research*, 8(6): 1-7.

